

# MATHEMATICS

## Chapter 9: Rational Numbers



## Rational Numbers

1. The numbers of the form  $\frac{x}{y}$ , where  $x$  and  $y$  are natural numbers, are known as fractions.
2. A number of the form  $\frac{p}{q}$  [ $q \neq 0$ ], where  $p$  and  $q$  are integers, is called a rational number.
3. Every integer is a rational number and every fraction is a rational number.
4. A rational number  $\frac{p}{q}$  is positive if  $p$  and  $q$  are either both positive or both negative.
5. A rational number  $\frac{p}{q}$  is negative if one of  $p$  and  $q$  is positive and the other one is negative.
6. Every positive rational number is greater than 0.
7. Every negative rational number is less than 0.
8. If  $\frac{p}{q}$  is a rational number and ' $m$ ' is a non-zero integer, then  $\frac{p}{q} \times \frac{p \times m}{q \times m}$ . Here,  $\frac{p}{q}$  and  $\frac{p \times m}{q \times m}$  are known as equivalent rational numbers.
9. If  $\frac{p}{q}$  is a rational number and ' $m$ ' is a common divisor of  $p$  and  $q$ , then  $\frac{p}{q} \times \frac{p \times m}{q \times m}$ . Here,  $\frac{p}{q}$  and  $\frac{p \times m}{q \times m}$  are known as equivalent rational numbers.
10. Two rational numbers are equivalent only when the product of numerator of the first and the denominator of the second is equal to the product of the denominator of the first and the numerator of the second.
11. A rational number  $\frac{p}{q}$  is said to be in standard form if  $q$  is positive and the integers  $p$  and  $q$  have no common divisors other than 1.
12. If there are two rational numbers with a common denominator then the one with the larger numerator is greater than the other.
13. Rational numbers with different denominators can be compared by first making their denominators same and then comparing their numerators.
14. There are infinite rational numbers between two rational numbers.
15. Two rational numbers with the same denominator can be added by adding their numerators, keeping the denominator same.

$$\frac{p}{q} \times \frac{r}{q} \times \frac{[p \times r]}{q}$$

16. Two rational numbers with different denominators are added by first taking the LCM of the two denominators and converting both the rational numbers to their equivalent forms having the LCM as the denominator.
17. While subtracting two rational numbers, we add the additive inverse of the rational number to be subtracted to the other rational number.

$$\frac{p}{q} \times \frac{r}{s} \times \frac{p}{q} \times (\text{additive inverse of } \frac{r}{s})$$

18. To multiply two rational numbers, we multiply their numerators and denominators separately, and write the product as  $\frac{\text{product of numerators}}{\text{product of denominators}}$
19. Reciprocal of  $\frac{p}{q}$  is  $\frac{q}{p}$ .
20. To divide one rational number by the other non-zero rational number, we multiply the first rational number by the reciprocal of the other.

## Rational Numbers

A rational number is defined as a number that can be expressed in the form

$$\frac{p}{q}$$

, where  $p$  and  $q$  are integers and  $q \neq 0$ .

In our daily lives, we use some quantities which are not whole numbers but can be expressed in the form of

$$\frac{p}{q}$$

Hence, we need rational numbers.

## Equivalent Rational Numbers

By multiplying or dividing the numerator and denominator of a rational number by a same non zero integer, we obtain another rational number equivalent to the given rational number. These are called equivalent fractions.

$$\left[ \frac{1}{3} = \frac{1}{3} \times \frac{2}{2} = \frac{2}{6} \right]$$

∴

$$\frac{2}{6}$$

and

$$\frac{1}{3}$$

are equivalent fractions.

$$\frac{15}{25} = \frac{15 \div 5}{25 \div 5} = \frac{3}{5}$$

∴

$$\frac{15}{25}$$

and

$$\frac{3}{5}$$

are equivalent fractions.

### Rational Numbers in Standard Form

A rational number is said to be in the standard form if its denominator is a positive integer and the numerator and denominator have no common factor other than 1.

Example: Reduce

$$\frac{-4}{16}$$

Here, the H.C.F. of 4 and 16 is 4.

$$\Rightarrow \frac{-4}{16} = \frac{\frac{-4}{4}}{\frac{16}{4}} = \frac{-4}{16}$$

$$\frac{-1}{4}$$

is the standard form of

$$\frac{-4}{16}$$

### LCM

The least common multiple (LCM) of two numbers is the smallest number ( $\neq 0$ ) that is a multiple of both.

Example: LCM of 3 and 4 can be calculated as shown below:

Multiples of 3: 0, 3, 6, 9, 12, 15

Multiples of 4: 0, 4, 8, 12, 16

LCM of 3 and 4 is 12.

### Rational Numbers between Two Rational Numbers

There are unlimited number (infinite number) of rational numbers between any two rational numbers.

Example: List some of the rational numbers between  $-35$  and  $-13$ .

Solution: L.C.M. of 5 and 3 is 15.

⇒ The given equations can be written as

$$\frac{-9}{15}$$

and

$$\frac{-5}{15}$$

⇒  $-615$ ,  $-715$ ,  $-815$  are the rational numbers between  $-35$  and  $-13$ .

Note: These are only few of the rational numbers between  $-35$  and  $-13$ . There are infinite number of rational numbers between them. Following the same procedure, many more rational numbers can be inserted between them.

## Properties of Rational Numbers

- **Closure Property**

Sum, difference and product of two rationals is again a rational number. So, Rational numbers are closed under addition, subtraction, multiplication but **NOT** under division.

- **Commutativity Property**

For any two rational numbers  $a$  and  $b$   $a * b = b * a$ .

- Rational numbers are commutative under addition and multiplication but **NOT** under subtraction and division.

Example:  $\frac{1}{7} + \frac{3}{7} = \frac{4}{7}$  and  $\frac{3}{7} + \frac{1}{7} = \frac{4}{7}$   
 $\frac{2}{3} \times \frac{5}{6} = \frac{10}{18} = \frac{5}{9}$  and  $\frac{5}{6} \times \frac{2}{3} = \frac{5}{9}$   
 $\frac{1}{2} - \frac{3}{4} = -\frac{1}{4}$  but  $\frac{3}{4} - \frac{1}{2} = \frac{1}{4}$   
 $\frac{3}{7} \div \frac{5}{2} = \frac{6}{35}$  but  $\frac{5}{2} \div \frac{3}{7} = \frac{35}{6}$

- **Associative Property**

For any three rational numbers  $a, b$  and  $c$ ,  $(a * b) * c = a * (b * c)$ .

- Addition and multiplication are associative for rational numbers, but subtraction and division are **NOT** associative for rational numbers.

Example:  $\left(\frac{1}{5} + \frac{2}{7}\right) + \frac{1}{3} = \frac{86}{105}$  and  $\frac{1}{5} + \left(\frac{2}{7} + \frac{1}{3}\right) = \frac{86}{105}$   
 $\left(\frac{3}{8} \times \frac{1}{9}\right) \times \frac{5}{7} = \frac{15}{504}$  and  $\frac{3}{8} \times \left(\frac{1}{9} \times \frac{5}{7}\right) = \frac{15}{504}$   
 $\left(\frac{4}{9} - \frac{3}{2}\right) - \frac{1}{3} = \frac{93}{57}$  but  $\frac{4}{9} - \left(\frac{3}{2} - \frac{1}{3}\right) = \frac{39}{54}$   
 $\left(\frac{3}{5} \div \frac{2}{5}\right) \div \frac{2}{5} = \frac{15}{4}$  but  $\frac{3}{5} \div \left(\frac{2}{5} \div \frac{2}{5}\right) = \frac{3}{5}$

## Addition of Rational Numbers

- **Case 1:** Adding rational numbers with same denominators:

$$\begin{aligned} \text{Example : } & \frac{19}{5} + \frac{-7}{5} \\ & = \left(\frac{19-7}{5}\right) = \frac{12}{5} \end{aligned}$$

- **Case 2:** Adding rational numbers with different denominators:

$$\text{Example : } \frac{-3}{7} + \frac{2}{3}$$

LCM of 7 and 3 is 21

$$\text{So, } \frac{-3}{7} = \frac{-9}{21} \text{ and } \frac{2}{3} = \frac{14}{21}$$

$$\Rightarrow \frac{-9}{21} + \frac{14}{21} = \left(\frac{-9+14}{21}\right) = \frac{5}{21}$$

## Subtraction of Rational Numbers

- To subtract two rational numbers, add the additive inverse of the rational number that is being subtracted, to the other rational number.

- Example: Subtract  $\frac{2}{5}$  from  $\frac{7}{9}$ .

$$\begin{aligned} & \frac{7}{9} + \text{Additive Inverse of } \left(\frac{2}{5}\right) \\ &= \frac{7}{9} + \left(\frac{-2}{5}\right) \\ &= \left(\frac{35-18}{45}\right) \quad \{\because \text{LCM of 9 and 5 is 45}\} \\ &= \frac{17}{45} \end{aligned}$$

## Multiplication and Division of Rational Numbers

### Multiplication of Rational Numbers

- Case 1:** To multiply a rational number by a positive integer, multiply the numerator by that integer, keeping the denominator unchanged.

$$\frac{-3}{5} \times (7) = \frac{-3 \times 7}{5} = \frac{-21}{5}$$

- Case 2:** Steps to multiply one rational number by the other rational number:

**Step 1:** Multiply the numerators of the two rational numbers.

**Step 2:** Multiply the denominators of the two rational numbers.

**Step 3:** Write the product as

$$\begin{aligned} & \frac{\text{Product of Numerators}}{\text{Product of Denominators}} \\ &= \left(\frac{-5}{7}\right) \times \left(\frac{-9}{8}\right) = \frac{-5 \times (-9)}{7 \times 8} = \frac{45}{56} \end{aligned}$$

### Division of rational numbers

- To divide one rational number by the other rational numbers we multiply the rational number by the reciprocal of the other.

$$\begin{aligned} \text{Example: } & \frac{-2}{3} \div \frac{1}{7} \\ &= \frac{-2}{3} \times \text{Reciprocal of } \frac{1}{7} \\ &= \frac{-2}{3} \times 7 \quad \{\because \text{Reciprocal of } \frac{1}{7} = 7\} \\ &= \frac{-14}{3} \end{aligned}$$

## Negatives and Reciprocals

- Rational numbers are classified as positive and negative rational numbers.

(i) When both the numerator and denominator of a rational number are **positive integers or negative integers**, then it is a positive rational number.

Example:  $\frac{3}{5}$  is a positive rational number.  $\frac{-3}{-5} = \frac{3}{5}$  is also a positive rational number.

(ii) When either numerator or denominator of a rational number is a **negative integer**, it is a negative rational number.

Example:  $\frac{-3}{5} = -\frac{3}{5}$  is a negative rational number.  $\frac{3}{-5} = -\frac{3}{5}$  is also a negative rational number.

- If the product of two rational numbers is 1 then they are called **reciprocals** of each other.

Example :  $\frac{2}{3}$  is reciprocal of  $\frac{3}{2}$ , since  $\frac{2}{3} \times \frac{3}{2} = 1$

Note : The product of a rational number with its reciprocal is always 1.

### Additive Inverse of a Rational Number

- Additive Inverse of a rational number  $\frac{p}{q}$  is the number that, when added to  $\frac{p}{q}$ , yields zero.

Example: Additive Inverse of a rational number  $\frac{3}{5}$  is  $\frac{-3}{5}$  and additive inverse of  $\frac{-3}{5}$  is  $\frac{3}{5}$ .

Since  $\frac{3}{5} + \frac{-3}{5} = 0$

### Rational Numbers on a Number Line



- In order to represent a given rational number  $\frac{a}{n}$ , where  $a$  and  $n$  are integers, on the number line :


**Step 1:** Divide the distance between two consecutive integers into  $n$  parts.

For example : If we are given a rational number  $\frac{3}{4}$ , we divide the space between 0 and 1, 1 and 2 etc. into **four** parts

**Step 2:** Label the rational numbers till the range includes the number you need to mark

- The following figure shows how fractions  $\frac{1}{4}$ ,  $\frac{2}{4}$  and  $\frac{3}{4}$  are represented on a number line.
- Divide the portion from 0 to 1 on the number line into four parts.

Then each part represents  $\frac{1}{4}$ <sup>th</sup> portion of the whole.



## Comparison of Rational Numbers

- **Case 1:** To compare two negative rational numbers, ignore their negative signs and reverse the order.

Example: Which is greater:  $-\frac{3}{8}$  or  $-\frac{2}{7}$ ?

Compare  $\frac{3}{8}$  and  $\frac{2}{7}$ :  $\frac{3}{8} > \frac{2}{7}$


$\therefore -\frac{3}{8} < -\frac{2}{7}$

- **Case 2:** To compare a negative and a positive rational number, we consider that a negative rational number is to the left of zero whereas a positive rational number is to the right of zero on a number line. So, a negative rational number will always be less than a positive rational number.

Example: (i)  $-\frac{3}{11} < \frac{2}{5}$

(ii)  $-\frac{3}{8} < -\frac{2}{7}$

Class : 7th mathematics  
Chapter-9 : Rational Numbers



## Important Questions

### Multiple Choice Questions:

Question 1. The numerator of the rational number  $\frac{1}{100}$  is

- (a) 100
- (b) 1
- (c) 10
- (d) 99

Question 2. The denominator of the rational number  $\frac{4}{7}$  is

- (a) 7
- (b) 4
- (c) 3
- (d) 11

Question 3. The denominator of the rational number  $\frac{7}{13}$  is

- (a) 13
- (b) 7
- (c) 6
- (d) 91

Question 4. The numerator of the rational number  $-\frac{3}{4}$  is

- (a) -3
- (b) 3
- (c) 4
- (d) -4

Question 5. The numerator of the rational number  $-\frac{2}{9}$  is

- (a) -2
- (b) 2
- (c) -9
- (d) 9

Question 6. The denominator of the rational number  $\frac{5}{-3}$  is

- (a) 5
- (b) -3

(c) 3

(d) 8

Question 7. The denominator of the rational number  $\frac{3}{-7}$  is

(a) 7

(b) -7

(c) 3

(d) -3

Question 8. The numerator of the rational number  $\frac{-2}{-5}$  is

(a) 2

(b) -2

(c) 5

(d) -5

Question 9. the numerator of the rational number  $\frac{-5}{-3}$  is

(a) -5

(b) 5

(c) -3

(d) 3

Question 10. The denominator of the rational number  $\frac{-2}{-9}$  is

(a) -2

(b) 2

(c) 9

(d) -9

Question 11. The denominator of the rational number  $\frac{-6}{-5}$  is

(a) 6

(b) -6

(c) 5

(d) -5

Question 12. The denominator of the rational number  $\frac{-13}{-11}$  is

(a) -13

(b) 13

(c) 11

(d) -11

Question 13. The numerator of the rational number 0 is

(a) 0

(b) 1

(c) 2

(d) 3

Question 14. The denominator of the rational number 0 is

(a) 0

(b) 1

(c) -1

(d) any non-zero integer

Question 15. The numerator of a rational number 8 is

(a) 2

(b) 4

(c) 6

(d) 8

### Very Short Questions:

1. Find three rational numbers equivalent to each of the following rational numbers.

(i)  $\frac{-2}{5}$

(ii)  $\frac{3}{7}$

2. Reduce the following rational numbers in standard form.

(i)  $\frac{35}{-15}$

(ii)  $\frac{-36}{-216}$

3. Represent  $\frac{3}{2}$  and  $\frac{-3}{4}$  on number lines.

4. Which of the following rational numbers is greater?

(i)  $\frac{3}{4}, \frac{1}{2}$

(ii)  $-\frac{3}{2}, \frac{-3}{4}$

5. Find the sum of

(i)  $-4\frac{3}{4} + 2\frac{7}{12}$       (ii)  $\frac{9}{-12} + \frac{5}{8}$

6. Subtract:

$$(i) \frac{-5}{6} \text{ from } \frac{-7}{8} \quad (ii) 2\frac{1}{5} \text{ from } -3\frac{1}{6}$$

### Short Questions:

1. Find the product:

$$(i) 6\frac{2}{3} \times \left(-5\frac{1}{16}\right) \quad (ii) \left(-3\frac{1}{4}\right) \times \left(-2\frac{3}{4}\right)$$

2. If the product of two rational numbers is  $\frac{-9}{16}$  and one of them is  $\frac{-4}{15}$ , find the other number.

3. Arrange the following rational numbers in ascending order.

$$(i) -\frac{1}{3}, \frac{-4}{3}, \frac{-2}{9} \quad (ii) -\frac{2}{3}, \frac{4}{5}, \frac{6}{7}, -\frac{1}{6}$$

4. Insert five rational numbers between:

$$(i) \frac{-2}{3} \text{ and } -1 \quad (ii) -\frac{1}{2} \text{ and } \frac{-3}{2}$$

5. Evaluate the following:

$$\frac{-12}{-5} + \frac{7}{-3} + \frac{-5}{14} + \frac{22}{7}$$

6. Subtract the sum of  $\frac{-5}{6}$  and  $-1\frac{3}{5}$  from the sum  $2\frac{2}{3}$  and  $-6\frac{2}{5}$ .

### Long Questions:

1.

$$\text{Simplify: } \left(\frac{3}{7} \times \frac{-5}{8}\right) + \left(\frac{1}{3} \times \frac{5}{6}\right) + \left|\frac{-1}{2} - \frac{1}{5}\right|$$

2. Divide the sum of  $-2\frac{15}{17}$  and  $3\frac{5}{34}$  by their difference.

3. During a festival sale, the cost of an object is ₹ 870 on which 20% is off. The same object is available at other shops for ₹ 975 with a discount of  $6\frac{2}{3}\%$ . Which is a better deal and by how much?

4. Simplify:

$$21.5 \div 5 - \frac{1}{5} \text{ of } (20.5 - 5.5) + 0.5 \times 8.5$$

5. Simplify:

$$2.3 - [1.89 - \{3.6 - (2.7 - \overline{0.8 - 0.03})\}]$$

6.

If  $x = \frac{-4}{9}$ ,  $y = \frac{5}{12}$  and  $z = \frac{7}{18}$ , find the value of

$$x \div y - \left[ \frac{1}{xy} - y \left( \frac{2x}{y} + \frac{x}{2y} \right) - xyz \left( \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \right) \right].$$

### ANSWER KEY -

#### Multiple Choice Questions:

1. (b) 1
2. (a) 7
3. (a) 13
4. (a) -3
5. (a) -2
6. (b) -3
7. (b) -7
8. (b) -2
9. (a) -5
10. (d) -9
11. (d) -5
12. (d) -11
13. (a) 0
14. (d) -1
15. (d) 8

#### Very Short Answer:

1.

$$(i) \frac{-2}{5} = \frac{-2 \times 2}{5 \times 2} = \frac{-4}{10}$$

$$\frac{-2}{5} = \frac{-2 \times 3}{5 \times 3} = \frac{-6}{15}$$

$$\frac{-2}{5} = \frac{-2 \times 4}{5 \times 4} = \frac{-8}{20}$$



Hence, the required rational numbers are

$$\frac{-4}{10}, \frac{-6}{15} \text{ and } \frac{-8}{20}.$$

$$(ii) \frac{3}{7} = \frac{3 \times 2}{7 \times 2} = \frac{6}{14}$$

$$\frac{3}{7} = \frac{3 \times 3}{7 \times 3} = \frac{9}{21}$$

$$\frac{3}{7} = \frac{3 \times 4}{7 \times 4} = \frac{12}{28}$$

Hence, the required rational numbers are

$$\frac{6}{14}, \frac{9}{21} \text{ and } \frac{12}{28}.$$

2.

$$(i) \frac{35}{-15} = \frac{-35}{15} = \frac{-35 \div 5}{15 \div 5} = \frac{-7}{3}$$

[ $\because$  HCF of 35 and 15 = 5]

$$(ii) \frac{-36}{-216} = \frac{36}{216} = \frac{36 \div 36}{216 \div 36} = \frac{1}{6}$$


[ $\because$  HCF of 36 and 216 = 36]

3.

$$(i) \frac{3}{2}$$



$$(ii) -\frac{3}{4}$$



4.



(i) We have  $\frac{3}{4}, \frac{1}{2}$

LCM of 4 and 2 = 4

$$\therefore \frac{3}{4} = \frac{3 \times 1}{4 \times 1} = \frac{3}{4}$$

$$\frac{1}{2} = \frac{1 \times 2}{2 \times 2} = \frac{2}{4}$$

Since  $\frac{3}{4} > \frac{2}{4}$

$$\therefore \frac{3}{4} > \frac{1}{2}$$

(ii) We have  $\frac{-3}{2}, \frac{-3}{4}$

LCM of 2 and 4 = 4

$$\therefore \frac{-3}{2} = \frac{-3 \times 2}{2 \times 2} = \frac{-6}{4}$$

$$\frac{-3}{4} = \frac{-3 \times 1}{4 \times 1} = \frac{-3}{4}$$

Since  $\frac{-3}{4} > \frac{-6}{4}$

$$\therefore \frac{-3}{4} > \frac{-3}{2}$$

5. Find the sum of

(i)  $-4\frac{3}{4} + 2\frac{7}{12}$

(ii)  $\frac{9}{-12} + \frac{5}{8}$

6.

$$= \frac{-7}{8} - \left(-\frac{5}{6}\right) = \frac{-7}{8} + \frac{5}{6}$$

$$= \frac{-7 \times 3}{8 \times 3} + \frac{5 \times 4}{6 \times 4} \quad [\text{LCM of 8 and 6} = 24]$$

$$= \frac{-21}{24} + \frac{20}{24} = \frac{-21 + 20}{24} = \frac{-1}{24}$$

(ii)  $2\frac{1}{5}$  from  $-3\frac{1}{6} = -3\frac{1}{6} - 2\frac{1}{5} = \frac{-19}{6} - \frac{11}{5}$

$$= \frac{-19 \times 5}{6 \times 5} - \frac{11 \times 6}{5 \times 6} \quad [\text{LCM of 6 and 5} = 30]$$

$$= \frac{-95}{30} - \frac{66}{30} = \frac{-95 - 66}{30}$$

$$= \frac{-161}{30} = -5\frac{11}{30}$$

## Short Answer:

1.

$$\begin{aligned}
 (i) \quad 6\frac{2}{3} \times \left(-5\frac{1}{16}\right) &= \frac{20}{3} \times \left(-\frac{81}{16}\right) \\
 &\Rightarrow \frac{5}{1} \times \left(-\frac{27}{4}\right) \\
 &\quad \left[\frac{20}{16} = \frac{20 \div 4}{16 \div 4} = \frac{5}{4}, \frac{-81}{3} = \frac{-81 \div 3}{3 \div 3} = \frac{-27}{1}\right] \\
 &= \frac{-5 \times 27}{1 \times 4} = \frac{-135}{4} = -33\frac{3}{4}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad \left(-3\frac{1}{4}\right) \times \left(-2\frac{3}{4}\right) &= \left(-\frac{13}{4}\right) \times \left(-\frac{11}{4}\right) \\
 &= (-) \times (-) \times \left(\frac{13}{4} \times \frac{11}{4}\right) = \frac{13 \times 11}{4 \times 4} \\
 &= \frac{143}{16} = 8\frac{15}{16} \quad [\because (-) \times (-) = (+)]
 \end{aligned}$$

2. Let the required rational number be x.

$$\begin{aligned}
 \therefore x \times \left(\frac{-4}{15}\right) &= -\frac{9}{16} \\
 \Rightarrow x &= -\frac{9}{16} \div \left(\frac{-4}{15}\right) = -\frac{9}{16} \times \frac{15}{-4} \\
 &\quad \left[\text{Reciprocal of } -\frac{4}{15} = \frac{15}{-4}\right] \\
 &= \frac{-9 \times 15}{-16 \times 4} = \frac{135}{64} = 2\frac{7}{64}
 \end{aligned}$$

Hence, the required rational number =  $2\frac{7}{64}$ .

3.

(i) We have  $-\frac{1}{3}, \frac{-4}{3}, \frac{-2}{9}$

LCM of 3, 3 and 9 = 9

$$\therefore \frac{-1 \times 3}{3 \times 3}, \frac{-4 \times 3}{3 \times 3}, \frac{-2 \times 1}{9 \times 1}$$

[Converting denominators as same number]

$$\Rightarrow \frac{-3}{9}, \frac{-12}{9}, \frac{-2}{9}$$

Since  $\frac{-12}{9} < \frac{-3}{9} < \frac{-2}{9}$

$$\therefore \frac{-4}{3} < -\frac{1}{3} < \frac{-2}{9}$$

(ii) We have  $-\frac{2}{3}, \frac{4}{5}, \frac{6}{7}, -\frac{1}{6}$

LCM of 3, 5, 6 and 7 = 210

$$\therefore \frac{-2 \times 70}{3 \times 70}, \frac{4 \times 42}{5 \times 42}, \frac{6 \times 30}{7 \times 30}, \frac{-1 \times 35}{6 \times 35}$$

[Converting the denominators as same numbers]

$$\Rightarrow \frac{-140}{210}, \frac{168}{210}, \frac{180}{210}, \frac{-35}{210}$$

Since  $\frac{-140}{210} < \frac{-35}{210} < \frac{168}{210} < \frac{180}{210}$

$$\therefore \frac{-2}{3} < -\frac{1}{6} < \frac{4}{5} < \frac{6}{7}$$

4.

$$(i) \frac{-2}{3} \text{ and } -1 \Rightarrow \frac{-2}{3} \text{ and } \frac{-1}{1}$$

LCM of 3 and 1 = 3

$$\therefore \frac{-2 \times 1}{3 \times 1} = \frac{-2}{3} \text{ and } \frac{-1 \times 3}{1 \times 3} = \frac{-3}{3}$$

We know that there is no integer between -2 and -3.

$\therefore$  Multiplying and dividing by  $5 + 1 = 6$  to each of the rational numbers, we have

$$\frac{-2 \times 6}{3 \times 6} = \frac{-12}{18} \text{ and } \frac{-3 \times 6}{3 \times 6} = \frac{-18}{18}$$

Here, integers between -12 and -18 are -13, -14, -15, -16 and -17.

$\therefore$  The required rational numbers are

$$\frac{-13}{18}, \frac{-14}{18}, \frac{-15}{18}, \frac{-16}{18} \text{ and } \frac{-17}{18}$$

$$\text{i.e., } \frac{-13}{18}, \frac{-7}{9}, \frac{-5}{6}, \frac{-8}{9}, \frac{-17}{18}$$

$$(ii) -\frac{1}{2} \text{ and } \frac{-3}{2}$$

Since, the denominator are same and there is only one integer between -1 and -3.

$\therefore$  Multiplying and dividing by  $5 + 1 = 6$  to each of the rational numbers, we have

$$\frac{-1 \times 6}{2 \times 6} = \frac{-6}{12} \text{ and } \frac{-3 \times 6}{2 \times 6} = \frac{-18}{12}$$

Here, the integers between -6 and -18 are -7, -8, -9, -10, -11

$\therefore$  The required rational numbers are

$$\frac{-7}{12}, \frac{-8}{12}, \frac{-9}{12}, \frac{-10}{12}, \frac{-11}{12}$$

$$\text{i.e., } \frac{-7}{12}, \frac{-2}{3}, \frac{-3}{4}, \frac{-5}{6}, \frac{-11}{12}$$

5.

$$\frac{-12}{-5} + \frac{7}{-3} + \frac{-5}{14} + \frac{22}{7}$$

$$= \frac{12}{5} - \frac{7}{3} - \frac{5}{14} + \frac{22}{7}$$

[LCM of 5, 3, 14, 7 = 210]

$$\therefore \frac{12}{5} = \frac{12 \times 42}{5 \times 42} = \frac{504}{210}$$

$$\frac{-7}{3} = \frac{-7 \times 70}{3 \times 70} = \frac{-490}{210}$$

$$\frac{-5}{14} = \frac{-5 \times 15}{14 \times 15} = \frac{-75}{210}$$

$$\frac{22}{7} = \frac{22 \times 30}{7 \times 30} = \frac{660}{210}$$

$$\begin{aligned} \text{So, } \frac{12}{5} - \frac{7}{3} + \frac{-5}{14} + \frac{22}{7} &= \frac{504}{210} - \frac{490}{210} - \frac{75}{210} + \frac{660}{210} \\ &= \frac{504 - 490 - 75 + 660}{210} \\ &= \frac{1164 - 565}{210} \\ &= \frac{599}{210} = 2\frac{179}{210} \end{aligned}$$

6.

SHIVOM CLASSES  
8696608541

$$\begin{aligned}
 &\text{Sum of } \frac{-5}{6} \text{ and } -1\frac{3}{5} \\
 \Rightarrow &\frac{-5}{6} + \left(-1\frac{3}{5}\right) = \frac{-5}{6} - \frac{8}{5} \\
 &= \frac{-5 \times 5}{6 \times 5} - \frac{8 \times 6}{5 \times 6} \\
 &\quad \text{[LCM of 6 and 5 = 30]} \\
 &= \frac{-25}{30} - \frac{48}{30} \\
 &= \frac{-25 - 48}{30} = \frac{-73}{30}
 \end{aligned}$$

$$\begin{aligned}
 &\text{Sum of } 2\frac{2}{3} \text{ and } -6\frac{2}{5} \\
 \Rightarrow &2\frac{2}{3} + \left(-6\frac{2}{5}\right) = \frac{8}{3} - \frac{32}{5} \\
 &= \frac{8 \times 5}{3 \times 5} - \frac{32 \times 3}{5 \times 3} \quad \text{[LCM of 3 and 5 = 15]} \\
 &= \frac{40}{15} - \frac{96}{15} = \frac{40 - 96}{15} = \frac{-56}{15}
 \end{aligned}$$

$\therefore$  Required difference is  $\left(\frac{-56}{15}\right) - \left(\frac{-73}{30}\right)$

$$\begin{aligned}
 \frac{-56}{15} + \frac{73}{30} &= \frac{73}{30} - \frac{56}{15} = \frac{73 \times 1}{30 \times 1} - \frac{56 \times 2}{15 \times 2} \\
 &= \frac{73}{30} - \frac{112}{30} = \frac{73 - 112}{30} = \frac{-39}{30} = \frac{-13}{10}
 \end{aligned}$$

**Long Answer:**

1. We have

$$\begin{aligned}
& \left( \frac{3}{7} \times \frac{-5}{8} \right) \div \left( \frac{1}{3} \times \frac{5}{6} \right) + \left| \frac{-1}{2} - \frac{1}{5} \right| \\
&= \left( \frac{-3 \times 5}{7 \times 8} \right) \div \left( \frac{1 \times 5}{3 \times 6} \right) + \left| \frac{-5-2}{10} \right| \\
&= \frac{-15}{56} \div \frac{5}{18} + \left| \frac{-7}{10} \right| \\
&= \frac{-15}{56} \times \frac{18}{5} + \frac{7}{10} \\
&\quad [\because \text{absolute value of } -a = a] \\
&= \frac{-27}{28} + \frac{7}{10} = \frac{-27 \times 5 + 7 \times 14}{140} \\
&\quad [\text{LCM of 28 and 10} = 140] \\
&= \frac{-135 + 98}{140} = \frac{-37}{140}
\end{aligned}$$

2.

Given rational numbers are  $-2\frac{15}{17}$  and  $3\frac{5}{34}$

Sum of the given numbers

$$\begin{aligned}
&= -2\frac{15}{17} + 3\frac{5}{34} = \frac{-49}{17} + \frac{107}{34} \\
&= \frac{-49 \times 2}{17 \times 2} + \frac{107 \times 1}{34 \times 1} \quad [\text{LCM of 17 and 34} = 34] \\
&= \frac{-98}{34} + \frac{107}{34} = \frac{-98 + 107}{34} = \frac{9}{34}
\end{aligned}$$

Difference of the given numbers =  $3\frac{5}{34} - \left(-2\frac{15}{17}\right)$

$$\begin{aligned}
&= \frac{107}{34} + \frac{49}{17} = \frac{107 \times 1}{34 \times 1} + \frac{49 \times 2}{17 \times 2} = \frac{107}{34} + \frac{98}{34} \\
&= \frac{107 + 98}{34} = \frac{205}{34}
\end{aligned}$$

As per the question, we have

Sum of the numbers  $\div$  Difference of the numbers

$$\frac{9}{34} \div \frac{205}{34} = \frac{9}{34} \times \frac{34}{205} = \frac{9}{205}$$

Hence, the required division =  $\frac{9}{205}$ .

3. The cost of the object = ₹ 870

$$\text{Discount} = 20\% \text{ of } ₹ 870 = \frac{20}{100} \times 870 = ₹ 174$$

$$\text{Selling price} = ₹ 870 - ₹ 174 = ₹ 696$$

The same object is available at other shop = ₹ 975

$$\begin{aligned} \text{Discount} &= 6\frac{2}{3}\% \text{ of ₹ 975} \\ &= \frac{20}{3} \times \frac{1}{100} \times 975 = ₹ 65 \end{aligned}$$

Selling price = ₹ 975 – ₹ 65 = ₹ 910

Since ₹ 910 > ₹ 696

Hence, deal at first shop is better and by ₹ 910 – ₹ 696 = ₹ 214

4. Using BODMAS rule, we have

$$\begin{aligned} &21.5 \div 5 - \frac{1}{5} \text{ of } (20.5 - 5.5) + 0.5 \times 8.5 \\ &= 21.5 \div 5 - \frac{1}{5} \text{ of } 15 + 0.5 \times 8.5 \\ &= 21.5 \times \frac{1}{5} - \frac{1}{5} \times 15 + 0.5 \times 8.5 \\ &= 4.3 - 3 + 4.25 \\ &= 4.3 + 4.25 - 3 \\ &= 8.55 - 3 \\ &= 5.55 \end{aligned}$$

5. Using BODMAS rule, we have

$$\begin{aligned} &2.3 - [1.89 - \{3.6 - (2.7 - 0.77)\}] \\ &= 2.3 - [1.89 - \{3.6 - 1.93\}] \\ &= 2.3 - [1.89 - 1.67] \\ &= 2.3 - 0.22 \\ &= 2.08 \end{aligned}$$

6. Using BODMAS rule, we have



$$\begin{aligned}
 x \div y - \left[ \frac{1}{xy} - y \left( \frac{2x}{y} \times \frac{2y}{x} \right) - \cancel{xyz} \left( \frac{yz + zx + xy}{\cancel{xyz}} \right) \right] \\
 = x \div y - \left[ \frac{1}{xy} - y(4) - (yz + zx + xy) \right] \\
 = x \div y - \left[ \frac{1}{xy} - 4y - yz - zx - xy \right] \\
 = x \div y - \frac{1}{xy} + 4y + yz + zx + xy \\
 = \frac{x}{y} - \frac{1}{xy} + 4y + yz + zx + xy
 \end{aligned}$$

Putting  $x = \frac{-4}{9}$ ,  $y = \frac{5}{12}$  and  $z = \frac{7}{18}$ , we get

$$\begin{aligned}
 &= \frac{-4}{9} - \frac{1}{\left(\frac{-4}{9}\right)\left(\frac{5}{12}\right)} + 4\left(\frac{5}{12}\right) \\
 &\quad + \left(\frac{5}{12}\right)\left(\frac{7}{18}\right) + \left(\frac{7}{18}\right)\left(\frac{-4}{9}\right) + \left(\frac{-4}{9}\right)\left(\frac{5}{12}\right) \\
 &= \frac{-4}{9} \times \frac{12}{5} - \frac{1}{\left(\frac{-5}{27}\right)} + \frac{5}{3} + \frac{35}{216} - \frac{14}{81} - \frac{5}{27}
 \end{aligned}$$